

Microbial Isolates from Patients and their Antibiogram at the Tertiary care Burn Unit in Bangladesh

H BEGUM^a, M QUAMRUZZAMAN^b, M TALUKDAR^c

Summary

Infection and antibacterial resistance are important issue in severe burn. This prospective study was carried out in 112 patients who were enrolled from July 2007 to June 2008 in DMCH burn unit (only dedicated burn facility in Bangladesh with a mean annual admission of 869). The aim of this study were to investigate the profile of microorganism and resistance to antimicrobial agents; individuals who were admitted more than 5 days, with partial or full thickness burn developed clinical signs and symptoms of wound infection or pneumonia or septicaemia were included in this study. Nearly 50% of participants were aged 11-30 yrs, the most common type of burn was flame burn and females were the common victims. Bacterial isolates were found in 104 (92.85%) samples and eight (08) swabs

were sterile (7.14%). Pseudomonas species was the commonest pathogens (46.42%) followed by Proteus species (21.41%) and Klebsiella species (19.6%). Multiple organisms were found in 38 samples(33.92%). Antibiogram results obtained from ten antimicrobial agents demonstrated that Imipenem was the most effective agent, followed by amikacin and ceftazidime (92.3%, 52.8% and 38.46% sensitive respectively). Hundred percent (100%) of Proteus species were sensitive to imipenem, then amikacin 58.33%; they were highly resistant to cephalosporins (nearly 100%) and ciprofloxacin (83.33%). The resistance of Pseudomonas aeruginosa to anti-pseudomonas agents were exceptionally high. Newer drugs were found to be effective.

(J Bangladesh Coll Phys Surg 2011; 29: 62-66)

Introduction:

Burn wounds are never sterile, even in the presence of topical agents or systemic antibiotics.¹ Severe burn caused complex changes in homeostasis that can hardly be compared with other traumas or disorders and mortality is relatively common in the early phase. Infection is an important cause of mortality in burn, if the burn exceeds 40% of the total body surface area (TBSA). Wound infection will be one of the main complications². Sepsis is the leading cause of death in patient with large burns.³ 75% of all deaths following burn injury are related to infection⁴. Infection in the wound prolongs the healing process; treatment includes

rational antibiotic administration, removal of necrotic tissues, sufficient blood and oxygen supply to the wound and good nutritional support in burn victims.⁵ Individual hospital units will notice a change in their common pathogens over time. Infections with Pseudomonas organisms began to be seen in increasing numbers.⁶ From 11% to 30% of burns are contaminated by microorganism of the gastrointestinal tract, skin and upper respiratory system, including Pseudomonas aeruginosa, Staphylococcus aureus, Escherichia coli, Klebsiella spp, Enterococcus spp and Candida spp.⁷ The organism most often involved in wound infection particularly in the first week is Staphylococcus aureus.⁸ Infection with gram-negative organism is more evident after the 1st week. Pseudomonas organism is present on the wounds of approximately 25% of burn patients.¹ Enterococci and Candida albicans are now seen with increasing frequency, each being found in the wounds of about 50% of burn patients¹. The rate of nosocomial infections are higher in burn patients due to various factors like nature of burn injury itself, immunocompromised status of the patients, invasive diagnostic and therapeutic procedures and prolonged ICU stay. In addition, cross-infection results between

a. Dr. Hasina Begum, DA, FCPS (Anaesthesiology), Junior Consultant, 50 Bedded Burn Unit, DMCH.

b. Dr. Muhammad Quamruzzaman, FCPS (Surgery); MS (Plastic Surgery), Assistant Professor (Plastic Surgery), 50 Bedded Burn Unit, DMCH.

c. Dr. Moumita Talukdar, DA, MCPS (Anaesthesiology), Medical Officer, 50 Bedded Burn Unit, DMCH.

Address of Correspondence : Dr. Hasina Begum, DA, FCPS (Anaesthesiology), Junior Consultant, 50 Bedded Burn Unit, DMCH.

Received : 23 September, 2008 **Accepted :** 18 October, 2010

different burn patients due to overcrowding in burn wards. Antimicrobial resistance is a great problem in infectious disease. In burn units, because of the wide use of antibiotics and particularly the empirical administration of broad-spectrum antimicrobial, this problem is worse than in other hospital department.⁹ Complicating this high rate of infection is the fact that the spectrum of bacterial isolate varies with time and geographical area.¹⁰

The present prospective study was conducted to determine the bacteriological profile and the resistance pattern in the burn unit of DMCH, a tertiary referral center, Bangladesh, which forms the basis for modification of drug regimen strategy.

Material and Methods:

This prospective cross-sectional study involved 112 patients who were enrolled from July 2007 to June 2008 in burn unit, DMCH, which is the only dedicated burn facility in Bangladesh with a mean annual admission of about 869. It receives all severe burn cases and more than 90% of burn cases in Bangladesh.

Early excision and skin grafting is practiced in our unit as soon as the patients are surgically fit and repeated till all areas of the body are grafted. Superficial burns not requiring surgery are dressed with duoderm to permit moist wound healing. Extent and severity of burns are calculated by assessing total percentage of body surface area (TBSA) burnt. Individuals who were admitted more

than 5 days, with partial or full-thickness burn who developed clinical signs and symptoms of wound infection or pneumonia or septicaemia were included in this study.

Demographic and clinical data including gender, age, degree of burn, TBSA burnt, cause of burn and antibiotic therapy were collected for each participant.

According to the clinical status of the participants, appropriate samples including wound swab, urine and blood samples were taken. The first swab was obtained from deep areas of the burn before any cleaning and transferred to the laboratory by sterile test tube. Blood and urine samples were collected from individuals with signs and symptoms of septicaemia or pneumonia. Blood sampling was repeated three times.

Descriptive statistics were used for demographic and bacteriological profiles of each case; χ^2 testing compared the microorganisms sensitivity and resistance patterns to the examined antimicrobial; $p < 0.05$ was considered significant.

Result:

A total 24, 354 patients were attended in DMCH burn unit from July 2007 to June 2008, of which 2376 (198/month) acute burn patients were admitted; 1613 (67.88%) patients were discharged with or without some residual deformity; 547 (23.02%) patients were expired; 557(23.56%) patients leave the hospital with or without consent of the authority. (Table-I).

Table-I

<i>Burn & Plastic Surgery Unit</i>									
Burn patients hospital outcome and pattern :									
Month	Total Patients Attended	Total Patients attended				Total Patients admitted	Discharged (%)	DORB & Absconded	Death (%)
	Emergency & OPD	Flame Burn	Scald Burn	Electric Burn	Chemical Burn				
July'07	1496	90	90	65	06	95	97	17	24
August'07	1829	145	137	10	04	60	87	06	24
September'07	1487	132	86	39	04	72	104	14	31
October'07	1477	93	49	37	05	80	134	31	35
November'07	1731	85	87	74	02	245	122	21	34
December'07	1833	112	96	68	07	317	207	54	51
January'08	2774	90	90	65	06	260	136	105	83
February'08	3348	145	137	10	04	314	128	96	89
March'08	3351	132	86	39	04	274	114	65	62
April'08	1537	93	49	37	05	221	171	65	41
May'08	1552	89	47	40	03	219	161	41	43
June'08	1939	88	50	49	04	219	152	45	30
Total	24354	1294	1004	533	54	2376	1613 (67.88%)	550 (23.56%)	547

A total 112 people were included in this study. Their characteristics including gender, distribution of age, cause of burn, and TBSA burnt are presented in Table - 2. Nearly 50% of the participant were aged 11 to 30 years. The most common type of burn was flame burn and females were the common victims.

Bacterial isolate were found in 104 (92.85%) samples and only eight (08) swabs were sterile 7(14%). Pseudomonas species was the commonest pathogen isolated (46.42%) followed by proteus species (21.4%), Klebsiella species (19.6%), Providencia (19.6%), E.coli (14.2%), Acinetobacter (12.5%) and Staphylococcus aureus (7.1%). Multiple organism were found in 38 samples (33.92%), None of the 5 blood samples contained them. Bacterial isolate were found in only in

9 urine samples (8.03%) of which pseudomonas species was the commonest (3.57%), as shown in Table-III.

Antibiogram of Pseudomonas Aeruginosa and Proteus species to 10 antimicrobial agents including carbapenems (Imipenems), Cephalosporins, fluoroquinolones, aminoglycosides and netilmicin are presented in Table-4. The most effective agent against P. aeruginosa was Imipenem (92.30%). After that, Amikacin and ceftazidime were the most effective agents at 53.80% and 38.46% respectively. Most isolates of P.aeruginosa were resistant to ampicillin (100%), cephalexin (96.15%), cefixime (92.30%) and ceftriaxone (88.46), 100% bacterial isolates of Proteus spp. were sensitive to imipenem, then amikacin 58.33%; they are highly resistant to cephalosporins (nearly 100%) and ciprofloxacin (83.33%).

Table-II

<i>Characteristics of burn patients (n-112)</i>												
Age Of the Patients			Gender of the patients				Causes of Burn			TBSA % of burn		
Age	No	%	Male	Female	Type	Number	%	% of	No	%		
in yrs								burn				
0-10	26	19.6	No.	%	No.	%	Scald	30	26.78	<10	14	12.5
11-20	20	14.28	40	35.71	72	64.28	Flame	50	44.64	11-20	20	17.85
21-30	38	32.14					Electric	22	19.64	21-30	32	28.57
31-40	10	8.9					Gas Explosion	10	8.9	31-40	28	25
>40	18	16.07					Chemical	0	0	41-50	10	8.9
										>50	8	7.14

TBSA = Total Burn Surface Area

Table-III

<i>Microorganisms isolated (n=112)</i>										
	Sample	Positive	No growth	Microorganism isolated						
				Pseudo.	Kleb.	Proteus	E. coli	Staph.	Acinetobacter	Providencia
Wound swab	112	104	08	52	22	24	16	08	14	22
		92.85%	7.14%	(46.42%)	(19.6%)	(21.4%)	(14.2%)	(7.1%)	(12.5%)	(19.6%)
Blood	05	No growth								
Urine	112	09	103	4	1	2	2	-	-	-
		(8.03%)	(91.9%)	(3.57%)	(0.89%)	(1.78%)	(1.78%)	-	-	-

Combined growth — 38 (33.92%)

Table-4a*Antibiogram for Pseudomonas spp. (n=52).*

Antibiotic	Sensitive	Intermediate	Resistance
Imipenem	48 (92.30%)	-	4 (7.6%)
Ciprofloxacin	6 (11.3%)	-	46 (88.46%)
Amikacin	28 (53.8%)	2 (3.8%)	22 (42.33%)
Gentamycin	6 (11.53)	2 (3.8%)	44 (84.61%)
Ceftazidime	20 (38.46%)	2 (3.8%)	30 (57.69%)
Ceftriaxone	2 (3.8%)	4 (7.6%)	46 (88.46%)
Cefixime	4 (7.6%)	-	48 (92.30%)
Cephalexin	2 (3.8%)	-	50 (96.15%)
Ampicilin	-	-	52 (100%)
Netilmicin	16 (30.76%)	-	36 (69.23%)

Table-4b*Antibiogram for Proteus spp. (n=24).*

Antibiotic	Sensitive	Intermediate	Resistance
Imipenem	24 (100%)	-	-
Ciprofloxacin	-	4 (16.66%)	20 (83.33%)
Amikacin	14 (58.33%)	2 (8.3%)	8 (33.33%)
Gentamycin	4 (16.66%)	-	20 (83.33%)
Ceftazidime	8 (33.33%)	-	16 (66.66%)
Ceftriaxone	2 (8.3%)	2 (8.3%)	20 (83.33%)
Cefixime	-	-	24 (100%)
Cephalexin	-	-	24 (100%)
Ampicilin	-	-	24 (100%)
Netilmicin	8 (33.33%)	-	16 (66.66%)

Discussion:

Burn patients constitute a subset of patients particularly prone to infection. In the present study the most commonly isolated organisms from burn patients were *Pseudomonas* species followed by *Proteus* spp. and *Klebsiella* spp. and the high resistance rate of *P. aeruginosa* to common anti *Pseudomonas* agents. Compared with other studies, it was highly corresponds with the study of G. Khorasania et. al². Like our study Agnihotri et al¹¹ reported that 96% of swab were positive and *P. aeruginosa* was found to be the most common isolates (59%). These results contrast to the study of Mehta Manjula et al¹². there was high incidence of *Staph aureus* isolation next to *Pseudomonas*. They also found a changing trend in burn bacteriology, it was

decreased for *Pseudomonas* spp. *Staph. aureus* and *Proteus* to increase for *Klebsiella* species. The prevalence of *Staph. aureus* was very low in our study (only 7.1%). Another significant difference of our study results to the other studies is that the prevalence of *Acinetobacter* spp. is very low (12.5%). There was significant rise in the isolation rate of *Acinetobacter* spp. over the last five to eight years as stated by Sengupta et al¹³. *Acinetobacter* spp. are emerging as an important cause of nosocomial infection in burn units. There are a number of factors which may contribute to this increase like its presence as a normal skin commensal and its easy spread due to multi drug resistance in a hospital settings.

G Khorasani et al² found high prevalence of *Citrobacter freundii* in their study and they found *Citrobacter* in all clinical samples except blood.

The high prevalence of *P. aeruginosa* is in agreement with other studies^{2,11,12,13,14} and may be explained by the fact that this opportunistic microorganism grows mainly in moist body areas, such as burn wounds¹⁵ and also by prolonged hospital stay and the administration of broad spectrum antibiotics in burn cases. A burn represents a site susceptible to opportunistic colonisation. The situation of burn victims with *P. aeruginosa* infection is particularly problematic, since this organism is inherently resistant to many drugs and is able to acquire resistance to all effective antimicrobial drugs.¹⁶ The change in the pattern of bacterial resistance in the burn unit has importance both for clinical settings and epidemiological purpose. We saw a significantly high percentage of resistance among gram-negative bacilli to aminoglycosides like gentamicin and amikacin, ciprofloxacin, amoxicillin, cefotaxime' and ceftriaxone. This alarming trend was seen for both Enterobacteriaceae group and *Pseudomonas* spp. as seen in the study of Mehta et al¹². A similar report of multi-drug resistant gram-negative bacilli was also reported by Singh et al¹⁷. In comparison, imipenem and combination of drugs were found to be effective. This could be due to the reason that these are reserve drugs and used as last options for multi drug resistant bacteria in our hospital settings. Such high antimicrobial resistance is probably promoted due to selective pressure exerted by an bacteria due to numerous reasons like non adherence to hospital; antibiotic policy and excessive and indiscriminate use of broad spectrum antibiotics.

These multidrug resistant strains establish themselves in the hospital environment in areas like sinks, taps, railing, mattress, toilets and thereby spread from one patient to another. The implementation of a strict infection control strategy with a more rational use of antibiotics, including topical formulations and antimicrobial rotation, has been proposed to prevent the high incidence of multi-drug resistant strains of microorganism, particularly of *P. aeruginosa* in burn units¹⁸.

Conclusion:

Routine microbiological surveillance and careful in vitro testing prior to antibiotic use and strict adherence to hospital antibiotic policy may help in the prevention and treatment of multi-drug resistant pathogens in burn infection.

References :

- Jesse B. Hall, MD. Grocery A. Schmidt, MD. Lawrance D.H. Wood, MD. PH; et al. Burns:resuscitation phase (Day 2 to Day 6), Principle of Critical care; 2nd ed. 1998;1437-1442.
- G.Khorasania, E. Salehifarb and G Eslamib; Profile of microorganism and antimicrobial resistance as tertiary care referral burn center in Iran; Science Direct-Burn; Accepted 11 Dec.2007; Available online 2 April 2008.
- Edward s-Jones V, Greenwood JE, Manchester Burns Research Group. What's new in Burn microbiology? JamesLaing Memorial Prize Essay 2000. Burns 2003; 29(1): 15-24.
- Vindenes H, Bjerknes R. Microbial colonization of Large wounds. Burns 1995;21:575-9.
- P.G. Bowler, B.I. Duerden and D.G. Armstrong, Wound microbiology and associated approaches to wound management, Clin Microbiol Rev 2001; 14: 244-269.
- Krizek T. Local factors influencing incidence of wound sepsis.Symposium on antibiotic prophylaxis and therapy. Contemp Surg 1977;10: 45-50.
- K.M. Ramakrishnan, V. Jayaraman, K. Ramachandran and T. Mahdivanan, The management of anaerobic infection in extensive burns, Burns 1986, 12: 270-272.
- Muller C:Burns and the immune network. J Trauma 1979; 19: 880.
- J. Vrankova and V. Adamkova, Bacteriological monitoring after burn injury, Acta Chir Plast 2004; 46: 48-50.
- Ananthakrishanan AN, Kanungo R, Kumar K, Badrinath S. Detection of extended spectrum beta Lactamase producers among surgical wound infections and burn patients in JIPMER. Indian J Med Microbiol 2002; 18: 160-5.
- N. Agnihotri, V. Gupta and R.M. Joshi, Aerobic bacterial isolates from burn wound infections and their antibiograms - a five-year study, Burns 2004; 30: 241-243.
- Mehta M, Dutta P, Gupta V. Bacterial isolates from burn wound infections and their antibiograms: A eight-year study. Indian J Plast Surg 2007; 40: 25-8
- Sengupta S, Kumar P, Ciraj AM, Shivananda PG. Acinetobacter baumannii - an emerging nosocomial pathogen in the burns unit. Manipal, India. Burns 2001; 27: 140-4.
- A.R. Lari, H.B. Honar and R. Alaghebandan, Pseudomonas infection in Tohid burn center, Iran, Burns 1998; 24: 637-641.
- W. Song, K.M. Lee, H.J. Kang, D.H. Shin and D.K. Kim, Microbiologic aspects of predominant bacteria isolated from the burn patients in Korea, Burns 2001; 27: 136-139.
- D.M. Livermore, Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare?, Clin Infect Dis 2002; 34: 634-640.
- Singh NP, Goyal R, Manchanda V, Das S, Kaur Z, Talwar V. Changing Trends in bacteriology of burns in the burns units, Delhi, India. Burns 2003; 29: 129-32.
- B.A. Cunha, Strategies to control antibiotic resistance, Semin Respir Infect 2002; 17: 250-258.